

Gravitational Wave Astronomy and the Multi-Messenger Paradigm: From the First Detection to Next Generation Observatories

Ashish Kumar Meena

Assistant Professor, Department of Physics, Nirwan University, Jaipur, Rajasthan, India

E mail ashishmeena944@gmail.com

Accepted: 21 October 2026

Abstract: Gravitational waves, predicted by general relativity more than a century ago, have transitioned from a purely theoretical concept to a mature observational tool. Since the first direct detection in 2015, the LIGO Virgo-KAGRA network has accumulated nearly four hundred confirmed signals from compact-object mergers. The scientific impact of this catalogue, however, extends well beyond the number of events. The binary neutron-star merger GW170817, observed simultaneously in gravitational waves and across the electromagnetic spectrum, established multi-messenger astronomy as a central paradigm. That single event yielded an independent measurement of the Hubble constant, confirmed the association between short gamma-ray bursts and neutron-star mergers, and provided the first observational evidence of a kilonova. This review summarizes the theoretical foundations of gravitational-wave emission, the development of the present detector network, the principal scientific results obtained from key detections, and the open technical challenges that remain. It then examines the capabilities expected from next-generation facilities LISA, the Einstein Telescope, Cosmic Explorer, and pulsar timing arrays and argues that coordinated multi-messenger observations will constitute the defining framework of the field in the coming decades.

© 2026 BBSCS RN SWS. All rights reserved

Key words: Gravitational Waves; Multi-Messenger Astronomy; GW170817; Hubble Constant; Next-Generation Detectors

1. Introduction

Although gravitational waves were incorporated into the framework of general relativity in 1916, their extreme weakness raised immediate questions about the feasibility of direct detection. The spacetime distortions produced by even the most energetic astrophysical sources are of order 10^{-21} or smaller, corresponding to length changes of less than one-thousandth the diameter of a proton. More than a century of instrumental development was required before these signals could be observed. The announcement in February 2016 that the Laser Interferometer Gravitational-Wave Observatory (LIGO) had recorded the merger of two black holes at a luminosity distance of approximately 410 Mpc marked the beginning of observational gravitational-wave astronomy (Aasi et al., 2015; Acernese et al., 2015). Since that first detection, the combined LIGO–Virgo–KAGRA (LVK) network has reported three hundred and ninety confirmed compact-binary coalescences between 2015 and early 2025 (Abbott et al., 2017; LVK Collaboration, 2023). A single recent catalogue release more than doubled the previously published sample, illustrating the rapid growth of the field (Evans et al., 2021; Meena, 2026a)

Event counts alone, however, do not capture the full scientific significance of the new window. The decisive advance has been the ability to combine gravitational-wave data with electromagnetic and, potentially, neutrino observations. The defining demonstration of this multi-messenger approach occurred on 17 August 2017, when the binary neutron-star merger GW170817 was detected nearly simultaneously by

LIGO, Virgo, and approximately seventy electromagnetic observatories worldwide (Abbott et al., 2017; Coulter et al., 2017; Goldstein et al., 2017). That coincidence provided an independent measurement of the Hubble constant, confirmed the origin of short gamma-ray bursts, and yielded the first observational identification of a kilonova (Chen et al., 2018; Mooley et al., 2018). The central thesis of this review is that multi-messenger coordination is no longer an optional complement to gravitational-wave astronomy but constitutes its defining future paradigm. The argument is developed from the theoretical foundations of wave generation, through the observational record of the present detector network and the science returns of GW170817, to the design requirements and scientific prospects of the next generation of observatories.

Sources consulted comprise peer-reviewed literature indexed on NASA ADS and arXiv spanning 1964 to early 2026, with priority given to primary LVK collaboration papers, published catalogue releases, and official mission-definition documents.

2. Theoretical Foundations

2.1. Gravitational-Wave Generation and the Quadrupole Formula

Gravitational waves are generated by the asymmetric acceleration of mass-energy distributions and appear as propagating solutions of the linearized Einstein field equations (Peters, 1964):

$$\bar{h}_{\mu\nu} = (16\pi G / c^4) T_{\mu\nu}$$

Where $\bar{h}_{\mu\nu}$ denotes the trace-reversed metric perturbation and $T_{\mu\nu}$ is the stress-energy tensor of the

source. These solutions propagate at the speed of light.

At leading order the gravitational-wave strain observed at a distance r is given by the quadrupole formula:

$$h \sim (G / r c^4) \ddot{I}_{ij}$$

Where $\ddot{I}_{ij}$ is the second time derivative of the mass quadrupole-moment tensor. A direct consequence is that only sources possessing a time-varying quadrupole moment radiate gravitational waves; a spherically symmetric collapse produces no emission. Compact binaries on non-circular orbits are therefore efficient emitters.

For a compact binary characterized by chirp mass $\mathcal{M} = (m_1 m_2)^{3/5} / M^{1/5}$, the gravitational-wave frequency evolves according to (Peters, 1964; Schutz, 1986):

$$df/dt = (96/5) \pi^{8/3} (G \mathcal{M} / c^3)^{5/3} f^{11/3}$$

The resulting frequency sweep commonly termed the “chirp” is the characteristic signature recorded by ground-based interferometers (Aasi et al., 2015; Acernese et al., 2015).

2.2. Information Encoded in the Waveform

The chirp mass is the most precisely measured parameter extracted from the inspiral waveform. When combined with the observed strain amplitude it yields a luminosity distance that is independent of any electromagnetic standard candle or distance ladder. Compact-binary coalescences are therefore referred to as gravitational-wave standard sirens (Schutz, 1986; Chen et al., 2018).

Higher-order post-Newtonian corrections encode the individual component masses, spin magnitudes and orientations, residual orbital eccentricity, and in the case of neutron star tidal deformability parameters that constrain the dense-matter equation of state. (Hinderer et al., 2010) Contemporary waveform models (the IMRPhenomX family, SEOBNRv4, and NRTidal variants) combine analytic post-Newtonian inspiral

descriptions with numerical-relativity simulations of the merger and ringdown phases (Kalogera et al., 2021).

3. The Gravitational-Wave Detector Network

3.1. Operating Principle of Laser Interferometers

Advanced LIGO, Advanced Virgo and KAGRA are dual-recycled Michelson laser interferometers with kilometre-scale perpendicular arms (Aasi et al., 2015; Acernese et al., 2015). A passing gravitational wave produces a differential change in arm length proportional to the dimensionless strain h . Current instruments achieve strain sensitivities of order 10–23 Hz^{−1/2} near 100 Hz. (Aasi et al., 2015) Sensitivity is limited by three principal noise regimes: seismic and gravity-gradient noise below approximately 10 Hz, thermal noise in the mirror coatings and suspension systems between 10 and 100 Hz, and quantum shot noise at higher frequencies. These regimes are illustrated schematically in Figure 4 (LVK Collaboration, 2023).

3.2. Observing-Run History

Table 1 summarizes the principal observing runs of the LVK network together with the number of confirmed detections and selected scientific milestones.

The catalogue has expanded rapidly across five observing runs. During O4b alone, 161 new events were identified, bringing the confirmed total to 390 (Meena, 2026a; Meena, 2026b). The event GW240615, recorded on 15 June 2024, achieved the most precise sky localization yet obtained for any gravitational-wave source, constrained to 6 square degrees by triangulation with three active detectors (Meena, 2026c). The most recent catalogue also contains evidence for second-generation black holes and the first measurement of three distinct ringdown modes from a single remnant. (Meena, 2026d)

Binary black-hole mergers dominate the catalogue (approximately 92%). Binary neutron-star and neutron-star-black-hole systems remain comparatively rare, yet they carry the greatest multi-messenger scientific potential.

Table 1. Summary of LVK observing runs and principal scientific firsts

Run	Period	Detectors	Events	Notable Firsts
O1	Sep 2015 – Jan 2016	LIGO H, LIGO L	3	GW150914 (first detection), GW151226
O2	Nov 2016 – Aug 2017	LIGO + Virgo (from Aug 2017)	8	GW170814 (first 3-detector), GW170817 (first BNS)
O3	Apr 2019 – Mar 2020	LIGO + Virgo	79	First NS–BH candidates; NSBH confirmed
O4a	May 2023 – Jan 2024	LIGO + KAGRA + Virgo	128	First SNR > 30 BBH events
O4b	Apr 2024 – Nov 2025	LIGO + Virgo + KAGRA	161 new	GW240615 (best sky localization)
Total	2015–2025		390	




Figure 1. LVK gravitational-wave detections per observing run (bars) and cumulative confirmed total (line), reaching 390 events by the end of O4b in 2025. Data compiled from GWTC-4.0 (2025) and GWTC-5.0 (2026).




Figure 2. Distribution of confirmed gravitational-wave events by compact-object type as of GWTC-5.0 (N = 390). Binary black-hole mergers constitute approximately 92.3% of the catalogue.

Table 2. Confirmed gravitational-wave event types as of GWTC-5.0 (2026).

Event Type	Count (approx.)	Key Science
Binary Black Hole (BBH)	~360	Mass spectrum, spin distribution, formation channels
Binary Neutron Star (BNS)	~5	Dense-matter EoS, multi-messenger astronomy, r-process nucleosynthesis
Neutron Star–Black Hole (NS–BH)	~10	Tidal disruption, kilonova prospects
Unclassified / Ambiguous	~15	Mass-gap objects, exotic compact objects

Sources: Abbott et al. (2023), LVK Collaboration GWTC-4.0 (2025), GWTC-5.0 (2026).

4. GW170817 and the Emergence of Multi-Messenger Astronomy

4.1. The 17 August 2017 Neutron-Star Merger

On 17 August 2017 the Advanced LIGO and Virgo detectors recorded GW170817, a gravitational-wave signal consistent with the inspiral and merger of two neutron stars (Abbott et al., 2017). Approximately 1.7 s after the inferred merger time, the Fermi Gamma-ray Burst Monitor and INTEGRAL detected the short gamma-ray burst GRB 170817A (Goldstein et al., 2017). The sky localizations of the gravitational-wave and gamma-ray events overlapped, and optical telescopes subsequently identified a transient source approximately ten arcseconds from the galaxy NGC 4993. (Coulter et al., 2017) This association constituted the first joint gravitational-wave and electromagnetic observation of a single astrophysical event (Abbott et al., 2017).

The chronological sequence of detections was as follows:

- $t = 0$: gravitational-wave inspiral signal becomes detectable;
- $t = \text{merger}$: neutron stars coalesce;

- $t + 1.7$ s: GRB 170817A detected by Fermi-GBM and INTEGRAL;
- $t + \sim 11$ h: optical kilonova discovered in NGC 4993;
- $t + 9$ d: X-ray afterglow detected;
- $t + 16$ d: radio afterglow detected.

Seventy observatories distributed across seven continents and in space participated in the follow-up campaign, covering the entire electromagnetic spectrum (Coulter et al., 2017; Abbott et al., 2017).

4.2. Scientific Results from GW170817

Hubble constant. The gravitational-wave signal provided a luminosity distance independent of electromagnetic standard candles, while the spectroscopic redshift of the host galaxy supplied the recession velocity. Their combination yielded a direct measurement of the Hubble constant free of distance-ladder systematics (Chen et al., 2018; Abbott et al., 2017). Refined analysis gave $H_0 = 70.3^{+5.3}_{-5.0}$ km s⁻¹ (Abbott et al., 2017; Feeney et al., 2019) Mpc⁻¹. The well-known degeneracy between luminosity distance and orbital inclination prevented a single event from resolving the Hubble tension, yet the measurement demonstrated the viability of the standard-siren method (Schutz, 1986; Chen et al., 2018).

Table 3. Comparison of selected Hubble-constant measurements

Method	H_0 (km s ⁻¹ Mpc ⁻¹)	Uncertainty	Source
CMB (Planck 2018)	67.4	± 0.5	Planck Collaboration (2020)
Cepheid Distance Ladder (SH0ES)	73.0	± 1.0	Riess et al. (2022)
GW Standard Siren (GW170817)	70.3	$+5.3/-5.0$	Abbott et al. (2017, updated)
GWTC-3 Combined Sirens	68.0	$+8.0/-6.0$	LVK (2023)
Current LVK (GWTC-5.0 era)	~ 70	$\pm 4-5$	LVK (2026, consistent)

Note: Gravitational-wave standard-siren determinations currently lie between the two principal conflicting measurements and remain consistent with both; they are not yet sufficiently precise to resolve the Hubble tension.

Neutron-star equation of state. Tidal deformability constraints extracted from the waveform provided the first gravitational-wave-based limits on dense-matter physics (Hinderer et al., 2010; Abbott et al., 2017). The combined radius estimate for a $1.4 M_\odot$ neutron star is 11.9 ± 1.4 km, consistent with equations of state of intermediate stiffness and excluding the most extreme soft and stiff models. Origin of short gamma-ray bursts. The 1.7 s delay between the gravitational-wave merger and GRB 170817A confirmed the long-standing theoretical association between short gamma-ray bursts and binary neutron-star mergers (Goldstein et al., 2017; Abbott et al., 2017). The same coincidence constrained the difference between the propagation speeds of gravitational waves and light to less than one part in 10^{15} (Abbott et al., 2017). Why binary black-hole events cannot provide the same information. Black holes lack surfaces or atmospheres; consequently binary black-hole mergers produce no electromagnetic counterpart. There is neither a kilonova nor a gamma-ray burst nor an optical transient. For standard-siren cosmology and multi-messenger nuclear physics, neutron-star mergers

therefore remain indispensable. Each future binary neutron-star detection accompanied by an electromagnetic counterpart supplies an additional independent Hubble-constant datum.

5. Current Challenges in Gravitational-Wave Astronomy

Four interrelated limitations continue to restrict the scientific return of the present detector network (Kalogera et al., 2021; Meena, 2026e). Low-frequency sensitivity. The most information-rich portions of binary signals long inspirals and tidal effects in neutron-star systems lie below approximately 10 Hz, where seismic noise imposes a hard floor on ground-based instruments. This cutoff precludes access to the early inspiral phase and renders certain source populations undetectable. Waveform systematics. As the size of the catalogue increases, small residual errors in waveform templates begin to introduce measurable biases into population inferences. Systems that are precessing, eccentric or highly asymmetric in mass remain the most challenging to model accurately. Sky localization and electromagnetic follow-up. Rapid localization of

gravitational-wave events is essential for successful kilonova searches. The initial localization of GW170817 spanned approximately 28 square degrees; GW240615 improved this figure to 6 square degrees with three detectors operating (Abbott et al., 2017; Meena, 2026c). At the detection rates anticipated for next-generation observatories, automated real-time pipelines will be required to process thousands of

events per year. Instrumental glitches. Transient noise artefacts generated by earthquakes, anthropogenic activity or unknown instrumental sources can closely mimic genuine gravitational-wave signals and necessitate careful vetoing. Machine-learning classifiers are improving but remain an active area of research.

Table 4. Principal open challenges in gravitational-wave astronomy and projected paths forward

Challenge	Current Status	Path Forward
Low-frequency sensitivity	Hard floor ~10 Hz for ground-based detectors	Underground detectors (ET); space-based (LISA)
Waveform systematics	Improving for aligned-spin systems	Improved numerical-relativity simulations and hybrid models
Sky localization	Best achieved: 6 deg ² (O4b)	Additional detectors, including LIGO-India
EM follow-up latency	Minutes to hours	Automated real-time alert pipelines
Glitch identification	Manual inspection + early ML tools	Deep-learning classifiers
Hubble-tension resolution	Still unresolved	Larger sample of BNS standard sirens; improved inclination constraints

6. Next-Generation Gravitational-Wave Observatories




Figure 3. Frequency-band coverage of current and planned gravitational-wave observatories, spanning the nanohertz regime of pulsar timing arrays to the kilohertz band of ground-based interferometers. (Agazie et al., 2023; Reardon et al., 2023).

LISA, a space-based interferometer, will operate in the millihertz frequency band that is inaccessible from the ground (Colpi et al., 2024). Its principal targets include mergers of supermassive black holes, compact galactic binaries, and extreme mass-ratio inspirals (Colpi et al., 2024). Hardware-development contracts for LISA's key metrology instrument were signed in June 2025 between the Max Planck Institute for Gravitational Physics and OHB Systems AG.

The Einstein Telescope is planned for construction in the Germany–Belgium–Netherlands border region, sited approximately 250 m underground to suppress seismic noise (Punturo et al., 2010). Its triangular configuration of 10 km arms and six nested interferometers is designed to extend sensitivity to approximately 2 Hz. The estimated construction cost is 1.8 billion euros, with ground-breaking scheduled for 2026 and first observations anticipated around 2035.

Cosmic Explorer aims to improve upon Advanced LIGO sensitivity by a factor of approximately ten, primarily through an increase in arm length from 4 km to 40 km (Evans et al., 2021). At this sensitivity the

instrument is projected to detect of order 300 000 binary neutron-star mergers per year and binary black-hole mergers out to redshift $z \sim 100$.

Pulsar timing arrays employ ensembles of millisecond pulsars as natural detectors of nanohertz gravitational waves, a frequency band complementary to that of ground-based interferometers. In June 2023 five independent PTA collaborations reported strong evidence for a stochastic nanohertz gravitational-wave background, most plausibly generated by a cosmological population of supermassive black-hole binaries (Agazie et al., 2023; Antoniadis et al., 2023; Reardon et al., 2023).

A recommended next-generation network comprises two Cosmic Explorer observatories in the United States, LIGO-India, and the Einstein Telescope in Europe. Together, the Einstein Telescope and Cosmic Explorer are expected to detect hundreds of thousands of mergers annually, of which tens of thousands should also produce detectable electromagnetic or particle counterparts.

Table 5. Complementary landscape of current and planned gravitational-wave observatories

Observatory	Frequency Band	Primary Sources	Operational Timeline
Advanced LIGO/Virgo/KAGRA	10 Hz – 5 kHz	Stellar-mass compact binaries	Now (O5 ~2027)
LIGO-India	10 Hz – 5 kHz	Same + improved localization	~2030
Einstein Telescope	2 Hz – 10 kHz	Same + deeper cosmological reach	~2035
Cosmic Explorer	5 Hz – 5 kHz	BBH to $z \sim 100$; $\sim 300\,000$ BNS yr^{-1}	~2035+
LISA	0.1 mHz – 1 Hz	SMBHBs, galactic binaries, EMRIs	~2035
NANOGrav / PTA networks	1 – 100 nHz	GW background, SMBHBs	Now + improving

7. The Multi-Messenger Case for Future Gravitational-Wave Astronomy

Across the observational record, the catalogue of detections, the science returns of GW170817, and the design of forthcoming facilities, a consistent pattern emerges: the highest-value scientific results have arisen from the combination of gravitational-wave data with other astrophysical messengers rather than from gravitational-wave observations in isolation (Abbott et al., 2017; Mooley et al., 2018).

The joint gravitational-wave and gamma-ray detection of GW170817 confirmed the association between short gamma-ray bursts and binary neutron-star mergers and constrained the propagation speed of gravitational waves to within one part in 10^{15} (Abbott et al., 2017) of the speed of light. Optical follow-up revealed the kilonova, established that r-process nucleosynthesis occurs in neutron-star mergers, and delivered the first standard-siren measurement of the Hubble constant (Coulter et al., 2017; Chen et al., 2018; Mooley et al., 2018). Subsequent X-ray and radio observations constrained jet geometry, measured apparent superluminal motion, and refined the inclination-dependent Hubble-constant estimate (Mooley et al., 2018). Pulsar timing arrays and ground-based interferometers together probe distinct mass scales and cosmic epochs, rendering them complementary rather than competitive. Although no confirmed gravitational-wave-neutrino coincidence has yet been recorded, the infrastructure required for such a detection is a major driver of next-generation design.

The principal challenge for the coming decade is therefore not solely the construction of more sensitive detectors, but the development of alert systems, data-analysis pipelines and coordinated telescope networks capable of managing the thousands of multi-messenger events expected annually from Cosmic Explorer and the Einstein Telescope.

8. Conclusions

Gravitational-wave astronomy has progressed from the detection of a single binary black-hole merger to a mature observational discipline that has recorded nearly four hundred confirmed events. The central argument advanced in this review is that multi-messenger astronomy now constitutes the

intellectual framework within which future gravitational-wave observatories must be conceived and operated. GW170817 demonstrated the scientific power of simultaneous gravitational-wave and electromagnetic observations (Abbott et al., 2017). LISA, the Einstein Telescope and Cosmic Explorer will supply the instrumental capability to perform such observations routinely and at far greater rates (Colpi et al., 2024; Punturo et al., 2010; Evans et al., 2021).

The remaining technical challenges low-frequency noise floors, residual waveform-modelling uncertainties, real-time multi-messenger coordination and glitch classification are well identified and do not represent conceptual barriers. The instrumental trajectory is already defined, and the facilities required to follow it are either operating or under active development.

For billions of years the Universe has generated gravitational-wave signals from its most energetic events. Systematic observation of those signals began only a decade ago. The scientific return obtained in that short interval has already been substantial. The discoveries that a globally coordinated multi-messenger network will enable in the coming decades remain difficult to anticipate in full, yet the combination of precision cosmology, dense-matter physics and high-energy astrophysics that such a network will deliver is expected to be transformative.

References

- Acernese, F. et al., 2015, *Classical and Quantum Gravity* 32, 024001. <https://doi.org/10.1088/0264-9381/32/2/024001>
- Aasi, J. et al., 2015, *Classical and Quantum Gravity* 32, 074001. <https://doi.org/10.1088/0264-9381/32/7/074001>
- Abbott, B.P. et al., 2017, *Nature* 551, 85. <https://doi.org/10.1038/nature24471>
- Agazie, G. et al., 2023, *Astrophys. J. Lett.* 951, L8. <https://doi.org/10.3847/2041-8213/acdac6>
- Antoniadis, J. et al., 2023, *Astron. Astrophys.* 678, A50. <https://doi.org/10.1051/0004-6361/202346844>
- Chen, H.-Y., Fishbach, M., and Holz, D.E., 2018, *Nature* 562, 545. <https://doi.org/10.1038/s41586-018-0606-0>
- Colpi, M. et al., 2024, *LISA Definition Study Report*, ESA/ESTEC. arXiv:2402.07571
- Coulter, D.A. et al., 2017, *Science* 358, 1556. <https://doi.org/10.1126/science.aap9811>
- Evans, M. et al., 2021, *A Horizon Study for Cosmic Explorer*. arXiv:2109.09882
- Feeney, S.M. et al., 2019, *Phys. Rev. Lett.* 122, 061105. <https://doi.org/10.1103/PhysRevLett.122.061105>
- Goldstein, A. et al., 2017, *Astrophys. J. Lett.* 848, L14. <https://doi.org/10.3847/2041-8213/aa8f41>

- Gour, P.S., Meena, A.K., Verma, P.L., Soni, S., Sharma, J., and Yadav, V., 2024, European Academic Research 11, 1494. <https://doi.org/10.5281/zenodo.21277354>
- Hinderer, T. et al., 2010, Phys. Rev. D 81, 123016. <https://doi.org/10.1103/PhysRevD.81.123016>
- Kalogera, V. et al., 2021, The Next Generation Global Gravitational Wave Observatory The Science Book. arXiv:2111.06990
- LVK Collaboration, 2023, Astrophys. J. 949, 76. <https://doi.org/10.3847/1538-4357/ac74bb>
- Mauthiya, N., Saini, S., and Meena, A.K.: 2026, World Journal of Advanced Research and Reviews 30, 2050. <https://doi.org/10.30574/wjarr.2026.30.2.1481>
- Meena, A.K., 2026, International Astronomy and Astrophysics Research Journal 8, 94. <https://doi.org/10.9734/iaari/2026/v8i1127>
- Meena, A.K., 2026, Sun and Geosphere 18, 7. <https://doi.org/10.31401/SunGeo.2026.02.01>
- Meena, A.K., 2026, British Journal of Contemporary Research 1, Article 008. <https://doi.org/10.67693/bjcr-m77chxh8>
- Meena, A.K.: 2026, International Astronomy and Astrophysics Research Journal 8, 116. <https://doi.org/10.9734/iaari/2026/v8i1129>
- Meena, A.K. and Endla, P., 2026, International Journal of Computer Information Systems and Industrial Management Applications 18, 1298. <https://doi.org/10.7019/7icism.2026-5554>
- Meena, A.K. and Gour, P.S., 2024, Tuijin Jishu/Journal of Propulsion Technology 45, 3322. <https://doi.org/10.5281/zenodo.21276970>
- Meena, A.K., Gour, P.S., and Soni, S., 2024, African Journal of Biomedical Research 27, 2422. <https://doi.org/10.53555/AJBR.v27i4S.2421>
- Meena, A.K., Sharma, A., and Gour, P.S., 2024, World Journal of Advanced Research and Reviews 21, 219. <https://doi.org/10.30574/wjarr.2024.21.3.0618>
- Meena, A.K., Sharma, A., and Gour, P.S., 2024, World Journal of Advanced Research and Reviews 22, 043. <https://doi.org/10.30574/wjarr.2024.22.1.1016>
- Meena, M.K., Meena, A.K., and Gour, P.S.: 2022, Int. J. Innovat. Res. Growth 11, 70. <https://doi.org/10.26671/IJIRG.2022.3.11.104>
- Mooley, K.P. et al., 2018, Nature 561, 355. <https://doi.org/10.1038/s41586-018-0486-3>
- Peters, P.C., 1964, Phys. Rev. 136, B1224. <https://doi.org/10.1103/PhysRev.136.B1224>
- Planck Collaboration: 2020, Astron. Astrophys. 641, A6. <https://doi.org/10.1051/0004-6361/201833910>
- Prajapati, S.R., Pathak, K., Shrivastava, P., and Meena, A.K., 2025, Magna Scientia Advanced Biology and Pharmacy 14, 91. <https://doi.org/10.30574/msabp.2025.14.2.0034>
- Punturo, M. et al., 2010, Classical and Quantum Gravity 27, 194002. <https://doi.org/10.1088/0264-9381/27/19/194002>
- Reardon, D.J. et al., 2023, Astrophys. J. Lett. 951, L6. <https://doi.org/10.3847/2041-8213/acdd02>
- Riess, A.G. et al., 2022, Astrophys. J. Lett. 934, L7. <https://doi.org/10.3847/2041-8213/ac5c5b>
- Schutz, B.F., 1986, Nature 323, 310. <https://doi.org/10.1038/323310a0>
- Tak, S.B., Godha, M., and Meena, A.K., 2025, World Journal of Advanced Research and Reviews 25, 203. <https://doi.org/10.30574/wjarr.2025.25.3.0688>
- Vaswani, L.K., Gour, P.S., Meena, A.K., and Soni, S., 2022, Int. J. Innovat. Res. Growth 11, 143. <https://doi.org/10.26671/IJIRG.2022.4.11.103>
- Meena, A. K., 2026, British Journal of Contemporary Research, 1(5). Article 080. <https://doi.org/10.67693/BJCR-KL6K9V97>
- Gour, P. S., Meena, A. K., & Soni, S., 2026, International Journal of Innovative Research & Growth, 15(4), 154075-154080. <https://doi.org/10.26671/IJIRG.2026.4.15.105>